

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/395	A1	(11) International Publication Number: WO 96/39184 (43) International Publication Date: 12 December 1996 (12.12.96)
(21) International Application Number: PCT/US96/08028 (22) International Filing Date: 30 May 1996 (30.05.96) (30) Priority Data: 08/477,348 6 June 1995 (06.06.95) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). (72) Inventors: GRUNFELD, Carl; 299 Douglass Street, San Francisco, CA 94114 (US). FUNK, Janet; 60 Park Ridge #2, San Francisco, CA 94131 (US). FEINGOLD, Kenneth, R.; 214 Forbes, San Rafael, CA 94901 (US). (74) Agent: BERLINER, Robert; Robbins, Berliner & Carson, 5th floor, 201 N. Figueroa Street, Los Angeles, CA 90012-2628 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: THERAPEUTIC SEPSIS TREATMENT USING ANTAGONISTS TO PTHrP (57) Abstract <p>Methods and compositions are provided for the treatment or prophylaxis of systemic inflammatory response syndrome by administering an antagonist to parathyroid hormone-related protein, such as antibodies to PTHrP.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

THERAPEUTIC SEPSIS TREATMENT
USING ANTAGONISTS TO PTHrP

5 Field of the Invention

 This invention relates generally to a method of treating severe inflammatory conditions, such as systemic inflammatory response syndrome, which results from e.g., sepsis, and more particularly to administering an
10 antagonist to parathyroid hormone-related protein.

 This invention was made with Government support under Grant No. DK 47846 awarded by the National Institutes of Health and funds from the Research Service of the Department of Veterans' Affairs. The Government
15 has certain rights in this invention.

Background of the Invention

 Systematic inflammatory response syndrome is the designation recently established by a group of
20 researchers to describe related conditions resulting from, for example, sepsis, pancreatitis, multiple trauma such as injury to the brain, and tissue injury, such as laceration of the musculature, brain surgery, hemorrhagic shock, and immune-mediated organ injuries. A variety of
25 different approaches have been suggested for treating inflammatory conditions or septic (endotoxin) shock.

 U.S. Patent 5,308,834, issued May 3, 1994, inventors Scott et al., discloses a method said to prevent endotoxemia in a subject by administering an
30 amount of a leukocyte protein (BPI) effective to prevent endotoxemia in the subject.

 Among the therapeutic approaches, antibodies directed against endotoxin or its components have been evaluated for their utility in immunotherapy of sepsis.
35 Murine and human monoclonal antibodies directed against the core lipopolysaccharide of the endotoxin have been reported to exert protection during Gram-negative

bacterial sepsis in animals. Dunn, *Transplantation*, 45, 424-429 (1988). Antibodies directed against lipid A also have been reported to have a protective effect in humans. Jaspers et al., *Infection*, 15 Supp. 2, S89-95 (1987).

5 Antibodies to the J5 mutant of *E. coli* are reported to be protective against septic shock in animals and humans. Cohen et al., *Lancet*, 1, 8-11 (1987); Law and Marks, *J. Infect. Dis.*, 151, 988-994 (1985). Antibodies to endotoxin core glycolipid have been reported to prevent

10 the serious consequences of Gram-negative infections in surgical patients. Baumgartner et al., *Lancet*, 2, 59-63 (1985). In addition, human monoclonal antibodies to *P. aeruzinosa* exotoxin A and exoenzyme S have been described as useful for this purpose. U.S. Patent 4,677,070,

15 issued June 30, 1987.

Many of the toxic effects of endotoxin are mediated by cytokines, hormones, and other small molecules. Blockade of these various mediators has been used to treat sepsis. In animal models prior administration of

20 antibody directed against TNF was reported to protect from the lethal effects of endotoxin. Beutler et al., *Science*, 229, 869 (1985). Also, antibody blockade experiments were reported showing that various cytokines, such as TNF, are mediators of the lethal effects of endotoxin. Tracey et al., *Nature*, 330, 662 (1987); Ohlsson et

25 al., *Nature*, 348, 550 (1990); Heinzel, *J. Immunol.*, 145, 2920 (1990); Doherty et al., *J. Immunol.*, 149, 1666 (1992); Bernhagen et al., *Nature*, 365, 756 (1993). More recently, either polyclonal or monoclonal antibodies to a human tumor necrosis factor binding protein (TBP-I) have been described for

30 application in modulating the response to tumor necrosis factor, such as to suppress deleterious effects of this cytokine. U.S. Patent 5,359,037, issued October 25, 1994, inventors Wallach et al.

The treatment of endotoxemia or sepsis by

35 passive immunization with endotoxin neutralizing antibodies or cytokine antibodies is a relatively new approach. However, to date many of the approaches

suggested for sepsis treatment have not proven very efficacious. It is likely that treatment for septicemia in the future will combine a plurality of approaches, in view of the large cascade of pro-inflammatory cytokines
5 unleashed during the host response to infection.

Since morbidity and mortality associated with endotoxemia remains high, new adjunct therapies are being sought because septicemia remains the leading cause of death in intensive care units in the United States.

10

Summary of the Invention

In one aspect of the present invention, a method of treating a patient for a systemic inflammatory response syndrome comprises administering a pharma-
15 ceutically effective amount of a parathyroid hormone-related protein blocker to the patient. Thus, patients are treated for a systemic inflammatory response syndrome resulting from conditions such as sepsis.

Comparison between groups of inventively
20 treated and control animals showed that the inventive treatment provided a significant protective effect from an induced inflammatory response syndrome mortality, although the treatment delayed but, ultimately, did not prevent death. Thus, therapy in accordance with the
25 invention preferably includes an additional (or a plurality of) approach(es) to block the cascade of pro-inflammatory cytokines.

Brief Description of the Drawings

30 Figure 1 graphically illustrates the effect of goat anti-serum directed against a 34 amino acid fragment of PTHrP on mice that have been administered lethal doses of endotoxin. The dashed line represents similarly
35 treated mice to which were administered naive goat serum as control antibody;

Figure 2 is similar to Fig. 1, but illustrates rabbit anti-serum administrations; and

Figure 3 is similar to Fig. 2, but lethality data from five separate experiments were combined for analysis.

5 Detailed Description of the Preferred Embodiments

Parathyroid hormone-related protein (PTHrP) was isolated and identified in the late 1980s. The amino acid sequence of human, rat, mouse, and chicken PTHrPs are known to the art and are discussed by Martin et al.,
10 *Crit. Rev. Biochem. Mol. Biol.*, 26, pp. 377-395 (1991). PTHrP and PTH, one of the primary hormones responsible for calcium homeostasis, appear to be derived from a common ancestral gene. However, PTHrP is more highly conserved than PTH. Although PTHrP shares little primary
15 sequence homology with PTH, N-terminal segments of both peptides bind with similar affinity to PTH receptors present in kidney and bone, the classic target tissues for PTH action. The 1-13 region of PTHrP is 70% homologous with the corresponding region of PTH. The 14-
20 34 region of PTHrP shares no homology with the 14-34 region of PTH, although there is apparently a similar tertiary, 3-dimensional or steric homology between PTHrP and PTH in the 14-34 region.

In malignancy, high circulating levels of
25 tumor-derived PTHrP cause hypercalcemia by inappropriately stimulating bone resorption and renal calcium reabsorption via interaction with PTH/PTHrP receptors at these sites. In contrast, PTHrP is barely detectable in the circulation of normal individuals. However, PTHrP is
30 widely expressed in normal tissues, thus leading to the hypothesis that this peptide normally acts at its site of production in a paracrine or autocrine fashion. Consistent with this hypothesis, the recently cloned PTH/PTHrP receptor, in addition to being present in bone
35 and kidney, has also been found to be expressed in many of the same tissues that produce PTHrP.

While PTHrP has been defined by its relation to PTH, a hormone that follows the normal endocrine paradigm of localized production and distant action, we believe that PTHrP may in fact act more like a cytokine than a classic hormone and that its effect on bone and calcium metabolism may be but one of many important biological functions. We have found a therapeutic role for antibodies to PTHrP in significantly protecting from sepsis lethality.

Although the precise cause of death from sepsis is unknown, hypotension is a hallmark of lethal endotoxemia. Alterations in vascular hemodynamics are also thought to contribute to the multisystem organ failure that accompanies sepsis. In septic shock, systemic vascular resistance is low due to vasodilation and, although cardiac output increases in an attempt to maintain blood pressure, cardiac contractility is decreased.

The present invention is the therapeutic use of a PTHrP antagonist. By "PTHrP antagonist" is meant to include compounds that block PTHrP activity at the PTHrP receptor, and which include PTH or PTHrP fragments (e.g. fragments with the 3-34 or 7-34 amino acid residues of PTH or PTHrP and fragments in which one or more amino acid residues have been replaced with analogues), monoclonal or polyclonal antibodies to PTHrP, and non-peptide analogs that can be designed to mimic the effects of peptides such as the 3-34 or 7-34 amino acid residue fragments. The polyclonal or monoclonal antibodies may be raised in rabbits, mice, or other animals or tissue cultured cells or can be products of cells of human origin. They may also be produced of recombinant DNA technology either in a form identical to that of the native antibody or as chimeric molecules, constructed by recombination of antibody molecules of man and animal origins or in other forms chosen to make the antibodies most suitable for use in therapy. The replacement of

amino acid residues and the amide forms (at the C terminus) for analogues are known. Illustrative suitable analogues in accordance with the invention are where Nle is at the 8 and 18 positions and Tyr at the 34 of an amidated 3-34 parathyroid (bovine or human) fragment, where Tyr is at the 34 position of an amidated 7-34 (bovine) fragment, the amidated 7-34 fragment of PTHrP (human), where Leu is at 11 and D-Trp is at 12 of an amidated 7-34 PTHrP (human) fragment, where Asn is at 10, Leu at 11, and D-Trp at 12 of the 7-34 PTHrP fragment, and where D-Trp is at 12 and Tyr at 34 of an amidated 7-34 PTHrP (bovine) fragment.

In accordance with the inventive method, the PTHrP antagonist is administered prophylactically or therapeutically (that is, before, simultaneously with, or after) infection has set in. For example, when administering prophylactically, one particularly considers patients at risk such as those suffering from severe thermal burns, receiving immunosuppressive therapy, undergoing extensive surgical procedures, organ transplantation, or suffering other serious injuries or disease.

The PTHrP antagonist as therapeutic agent is administered to the patient by any suitable technique, preferably parenteral and, if desired intralesional. The specific method of administration will depend, e.g., on whether the administration is therapeutic or prophylactic. Thus, in view of the therapeutic urgency usually attending shock, the PTHrP antagonist may be intravenously infused at the same time as solutions used for initial volume expansion. Continuous infusion is preferred for administering peptides while bolus infusion may be used when administering antibodies. Prophylaxis is generally accomplished, e.g., by intramuscular or subcutaneous administration or other parenteral administration, including intraarterial and

intraperitoneal administration, preferably intravenous or intraperitoneal.

The PTHrP antagonist compositions to be used in the inventive therapy will be formulated and dosed in a fashion consistent with good medical practice taking into account the clinical condition of the individual patient, the cause of the septic shock, whether the PTHrP antagonist is used for therapy of shock or prophylaxis of incipient septic shock, the site of delivery of the PTHrP antagonist, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of the PTHrP antagonist administered parenterally per dose will be in the range of approximately 1 μ g/kg to 10 mg/kg of patient body weight once per day, although, as noted above, this will be subject to a great deal of therapeutic discretion. As earlier noted, where the PTHrP antagonist is a peptide or an analog of a peptide, then administration is preferably by continuous infusion. The key factor is selecting an appropriate dose and scheduling is the result obtained. Relatively higher doses may be needed initially for the treatment of profound shock, i.e., for patients in acute renal failure or respiratory distress, or having severely depressed blood pressure (mean arterial pressure below about 60 mm Hg).

For parental administration, the PTHrP antagonist is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a physiologically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed. Preferably the carrier is a parenteral carrier. Examples of such carrier vehicles include water, saline, Ringer's solution, dextrose solution, and

5% human serum albumin. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes. Generally, the carrier can contain minor amounts of additives such as substances that enhance isotonicity and chemical stability, e.g., buffers and preservatives, as well as low molecular weight (less than about 10 residues) polypeptides, proteins, amino acids, carbohydrates including glucose or dextrans, chelating agents such as EDTA, or other excipients. The PTHrP antagonist is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml at pH range 4 to 7.

A PTHrP antagonist for use in therapeutic administration must be sterile. This is readily accomplished by sterile filtration through (0.2 micron) membranes. The PTHrP antagonist selected ordinarily will be stored as an aqueous solution or can be lyophilized.

PTHrP antagonist therapy or prophylaxis is suitably combined (indeed, preferably combined) with other proposed or conventional therapies or prophylactic treatment for septic shock. For example, for treatment of burns, the PTHrP antagonist therapy may be delivered by separate means, simultaneously with and by the same administration route as other substances such as antibiotics, or anti-microbial agents that inhibit bacterial colonization of the burn wound surface. Other therapies that can be combined with PTHrP antagonist therapy include other cytokine antagonists, such as anti-TNF, antagonists of IL-1, and inhibitors of platelet activating factor. For example, Ohlsson et al., *supra*, have reported that a specific interleukin-1 receptor agonist reduces mortality from endotoxin shock while Doherty et al. and Heinzl, *supra*, discuss the use of anti-IFN- γ to reduce mortality from endotoxic shock. Further, the inventive method can be practiced in conjunction with primary therapeutic agents, for example, potent anti-microbial agents such as amino-glycosides

(such as amikacin, tobramycin netilmicin, and gentamicin), cephalosporin, related beta lactam agents such as moxalactam, carbopenems such as imipenem, monobactam agents such as aztreonam, ampicillin and
5 broad-spectrum penicillins (e.g., penicillinase-resistant penicillins, ureidopenicillins, or antipseudomonal penicillins).

It is known that the pathophysiologic consequences of Gram-negative sepsis are primarily
10 mediated by the release of bacterial endotoxin. Thus, endotoxin treatment of mice is used as an experimental model of gram negative sepsis. For example, in U.S. Patent 5,308,834, a group of rats was given a single, bolus injection of 0.5 mg/kg body weight bacterial
15 endotoxin in studying the effects of leukocyte protein ("BPI"), with the results said to support the use of BPI to reduce mortality due to sepsis. Similarly, U.S. Patent 5,055,447, issued October 8, 1991, inventors Palladino et al., used mice as a model in endotoxin
20 studies.

We have also used the injection of a near-lethal dose of endotoxin as a model for septic shock (a state of profound hypotension and multi-organ failure resulting from a systemic inflammatory response to
25 overwhelming infection by Gram-negative or other bacteria) in mice and rats as models of sepsis.

Thus, mice were passively immunized with antibody generated against the 1-34 fragment of PTHrP, which is a peptide fragment that is active at the
30 PTH/PTHrP receptor, prior to the administration of a lethal dose of endotoxin. Anti-PTHrP antibodies were raised both in goats and in rabbits by immunizing the animals with synthetic human PTHrP.

The immunoglobulin fraction of immune sera
35 (PTHrP antibody) or naive sera (control antibody) was partially purified by ammonium sulfate precipitation (33% saturation) using previously described techniques to

avoid the introduction of endotoxin contamination. The endotoxin content of PTHrP and control antibody solutions, determined by Limulus assay (sensitivity, 10 pg/ml), was below the levels required to alter sensitivity to subsequent (6 h) LPS challenge in mice.

Administration of goat PTHrP antibody 6 h prior to the injection of the endotoxin ("LPS") significantly protected mice from death when compared to the mortality rate (LD_{50}) seen in mice treated with goat control antibody ($p < 0.03$ by log rank analysis of Kaplan Meier curves) (Fig. 1). The protective effect of PTHrP antibody was confirmed in other studies utilizing antiserum raised in a different species (rabbit) against the same antigen. Passive immunization of mice with rabbit PTHrP antibody similarly protected mice from death caused by administration of LPS when compared to control mice treated with rabbit control antibody ($p < 0.004$ by log rank test of Kaplan Meier survival curves) (Fig. 2). The degree of protection from LPS lethality seen here with passive immunization against PTHrP is similar to that reported for passive immunization of mice against TNF, a major mediator of sepsis.

Comparison of survival curves for PTHrP antibody- versus control antibody-treated groups showed that PTHrP antibody provided a significant protective effect from LPS-induced death over the 3 to 4 day course of the experiments (Figs. 1 and 2), although the relative survival rates at the end of the observation periods were no different (as determined by Fisher's exact test). Thus, the PTHrP antibodies administered delayed but, ultimately, did not prevent death. Compilation of data from multiple experiments ($n=90$ mice) examining the ability of rabbit PTHrP antibody to protect from LPS-induced death confirms this conclusion. While overall survival was improved by pretreatment with rabbit PTHrP antibody ($p < .00005$ by log rank analysis), comparison of lethality rates at 12 h intervals following LPS

administration only showed a significant protective effect during the first 48 hours following LPS administration (Fig. 3).

In summary, these data show that PTHrP is effective in mediating at least some of the toxic effects of endotoxin, as evidenced by the ability of antibody directed against PTHrP to delay LPS-induced death.

Antibody blockade experiments, similar to those presented here, have previously shown that numerous other cytokines, such as TNF, IL-1, macrophage inhibitory factor (MIF) and IFN- γ , are also mediators of the lethal effects of endotoxin. The fact that passive immunization against PTHrP delays, but does not ultimately prevent, lethality from LPS is consistent with the hypothesis that PTHrP is one member of a larger cascade of pro-inflammatory cytokines that is unleashed during the host response to infection, and therapy in accordance with the invention will preferably include one or more approaches to block the cascade.

Turning to Fig. 1 and Fig. 2, the data shows the effect of goat (Fig. 1) or rabbit (Fig. 2) antiserum directed against PTHrP on LPS-induced lethality in mice. The details of the data are summarized by Figs. 1-3 are given by Examples 1-3.

EXAMPLE 1

Male C57BL/6 5-6 week old mice (n=9/group) were injected intraperitoneally (ip) with 200 μ l ammonium sulfate-precipitated goat antisera (PTHrP antibody, open triangle) (titre 1:8,000 by ELISA; LPS, undetectable) or ammonium sulfate-precipitated naive goat sera (control antibody, open circle) (LPS, undetectable) 6 h prior to the administration of 700 μ g 055:B5 LPS diluted in apyrogenic 0.9% saline. Animals, given access to chow and water *ad libitum*, were monitored for lethality for 96 h after LPS treatment. Statistical analysis of the

Kaplan Meier survival curves for the two groups (Statistica 4.1, StatSoft) using either a log rank test which gives equal weight to all points or the Peto & Peto Wilcoxon Test which weights earlier time points more heavily showed that pretreatment with PTHrP antibody protected mice from LPS-induced lethality ($p < 0.030$ and $p < 0.031$, respectively).

EXAMPLE 2

10

Mice ($n=8/\text{group}$) were injected with $100 \mu\text{l}$ ammonium sulfate precipitated rabbit antisera (open triangle) (titre 1:16,000; 6-fold concentrated by volume; $100 \text{ pg LPS}/100 \mu\text{l}$) or ammonium sulfate-precipitated naive rabbit serum (open circle) (5-fold concentrated by volume; $165 \text{ pg LPS}/100 \mu\text{l}$) 6 h prior to the administration of $700 \mu\text{g LPS}$ ip. Animals were then monitored as in (A). Statistical analysis of the Kaplan Meier survival curves for the two groups showed that pretreatment with PTHrP antibody protected mice from LPS-induced death ($p < 0.004$ by log rank test; $p < 0.003$ by Peto & Peto Wilcoxon Test). Results are representative of 5 separate experiments.

25

EXAMPLE 3

Turning to Fig. 3, the data shows the effect of rabbit antisera directed against PTHrP on LPS-induced lethality. Lethality data from 5 separate experiments, performed as described in Figs. 1 and 2, were combined for analysis. For a given experiment, C57BL/6 mice received an equal volume ($100\text{-}500 \mu\text{l}$) of PTHrP or control rabbit antibody. Since PTHrP antibody was ammonium sulfate-precipitated from different rabbit bleeds for different experiments, the volume of ammonium sulfate precipitated-PTHrP antibody used was adjusted so that an equivalent titer of PTHrP antibody was administered in

all experiments (e.g. 100 μ l of 1:16,000; 300 μ l of 1:5,500; or 500 μ l of 1:3,200). Control and PTHrP antibody solutions were matched for fold-concentration (4-6-fold concentrated relative to serum) and contained less than 165 pg LPS/volume injected. Six hours after antibody administration, mice from both treatment groups were injected with an equal amount of LPS (650-800 μ g LPS, corresponding to an LD₅₀ at t=48 h in control-antibody treated mice). Statistical analysis of Kaplan Meier survival curves (not shown here) for PTHrP (n=43) and control (n=47) antibody-treated groups showed that pretreatment with PTHrP antibody protected mice from LPS-induced death ($p < 0.00005$ by log rank analysis and $p < 0.000001$ by Peto & Peto Wilcoxon test). Comparison of lethality at 12 h intervals after LPS administration (from t=0-96 h) using Fisher's exact test (InStat 2.01, GraphPad Software, San Diego, CA) showed that PTHrP antibody significantly protected animals from LPS-induced lethality when compared to control antibody-treated mice at the early time points shown.

It is to be understood that while the invention has been described above in conjunction with preferred specific embodiments, the description and examples are intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims.

It is Claimed:

1. A method comprising:
administering to a patient suffering from risk
5 of systemic inflammatory response syndrome (SIRS) a
pharmaceutically effective amount of parathyroid hormone-
related protein (PTHrP) antagonist, the amount
administered being effective in reducing or delaying at
least some of the toxic effects of endotoxin or cytokine.
10
2. The method as in claim 1 wherein the
pharmaceutically effective amount is about 1 μ g/kg to 10
mg/kg of patient body weight.
- 15 3. The method as in claim 1 wherein the
administration is by continuous infusion or bolus
infusion.
4. The method as in claim 1 wherein the
20 administration is to a patient having a microbial
infection but not yet showing symptoms of septic shock.
5. The method as in claim 1 wherein the
antagonist is administered with an anti-microbial agent.
25
6. The method as in claim 1 wherein the PTHrP
antagonist administered includes monoclonal or polyclonal
antibodies to PTHrP.
- 30 7. The method as in claim 1 wherein the SIRS
is septic shock and the PTHrP antagonist administered
includes monoclonal or polyclonal antibodies to PTHrP.
8. The method as in claim 7 wherein the
35 administering is in conjunction with an additional
cytokine antagonist.

9. The method as in claim 1 wherein the PTHrP antagonist is administered with one or more of an anti-TNF agent, an IL-1 antagonist, and an inhibitor of platelet activating factor.

FIG. 1

1/1





FIG. 2




FIG. 3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/08028

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 39/395

US CL : 424/145.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/145.1, 158.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CA, MEDLINE

Search terms: systemic inflammatory response syndrome, SIRS, sepsis, septic, cytokine, antibody, PTHRP, parathyroid hormone

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DINARELLO et al. Anticytokine strategies in the treatment of the Systemic Inflammatory Response Syndrome. Journal of the American Medical Association. 14 April 1993, Vol. 269, No. 14, pages 1829-1835, especially abstract and pages 1829-1830.	1-9
Y	Chem. abstr., Vol. 118, No. 11, 15 March 1993 (Columbus, Ohio, USA), abstract 118: 100365w, JP 04-228089 A (EGUCHI, Y.) 18.08.92.	1-9
Y	FUNK et al. Endotoxin Increases Parathyroid Hormone-related Protein mRNA Levels in Mouse Spleen. Journal of Clinical Investigation. November 1993, Vol. 92, pages 2546-2552, especially abstract and pages 2546 and 2551.	1-9

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 26 JUNE 1996	Date of mailing of the international search report 12 JUL 1996
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>Prasad Murthy</i> PRASAD MURTHY Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/08028

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,055,447 A (PALLADINO ET AL) 08 October 1991, column 3, lines 53-58, column 4, lines 54-68 and column ,7 lines 51-68.	5, 8, 9